

Clinical Drug Testing: Common ICD-10-CM Sign and Symptom Codes

Listed below are the ICD-10-CM codes commonly received by Labcorp from ordering physicians for clinical drug testing performed at Labcorp and its MedTox Laboratories. Code descriptions were taken from the official code set, issued by the US Department of Health and Human Services, as updated effective October 1, 2024 through September 30, 2025. Depending on the nature of your practice, this list may be a useful reference tool; however, it is not a complete list. Please refer to an ICD-10-CM manual for a complete listing. Responsibility for the provision of correct diagnosis code(s) is with the ordering physician. The ordering physician must always determine, for the specific date of service, the appropriate diagnosis code(s) based on the patient's signs and symptoms. ICD-CM codes in effect at the time of service are required by third-party payers to support the medical necessity of the test(s) and/or profile(s) ordered. Physicians are asked to submit all medically appropriate ICD-CM codes when ordering laboratory testing.

Labcorp's provision of this information as a customer service does not suggest and is not meant to suggest that any of these codes should or should not be used on any given occasion. Labcorp makes no recommendation regarding the use of any particular diagnosis code(s). Diagnosis information should be reflected in the patient's medical record.

Symbol	Diagnosis	
	Code	Sign or Symptom
◀§	R79.9	Abnormal finding of blood chemistry, unspecified
	R74.8	Abnormal levels of other serum enzymes
§	R94.6	Abnormal results of thyroid functions studies
★§	J02.9	Acute pharyngitis, unspecified
▲§	M81.0	Age-related osteoporosis without current pathological fracture (adult age 15-124)
●§	F10.10	Alcohol abuse, uncomplicated
●§	F10.21	Alcohol dependence, in remission
●§	F10.20	Alcohol dependence, uncomplicated
§	D64.9	Anemia, unspecified
§	F50.019	Anorexia nervosa, restricting type, unspecified
	F41.9	Anxiety disorder, unspecified
	F90.2	Attention-deficit hyperactivity disorder, combined type
	F90.0	Attention-deficit hyperactivity disorder, predominantly inattentive type
	F90.9	Attention-deficit hyperactivity disorder, unspecified type
➤§	F84.0	Autistic disorder
§	F31.31	Bipolar disorder, current episode depressed, mild
§	F31.32	Bipolar disorder, current episode depressed, moderate
§	F31.5	Bipolar disorder, current episode depressed, severe, with psychotic features
§	F31.4	Bipolar disorder, current episode depressed, severe, without psychotic features
§	F31.2	Bipolar disorder, current episode manic severe with psychotic features
§	F31.12	Bipolar disorder, current episode manic without psychotic features, moderate
§	F31.62	Bipolar disorder, current episode mixed, moderate
§	F31.0	Bipolar disorder, current episodic hypomanic
§	F31.9	Bipolar disorder, unspecified
§	F31.81	Bipolar II disorder
§	F12.10	Cannabis abuse, uncomplicated
§	F12.20	Cannabis dependence, uncomplicated
	Z22.322	Carrier or suspected carrier of methicillin resistant staphylococcus aureus
§	M54.2	Cervicalgia
§	R53.82	Chronic fatigue, unspecified
▶§	G89.4	Chronic pain syndrome
✦§	B18.2	Chronic viral hepatitis C
§	F14.10	Cocaine abuse, uncomplicated
§	F14.20	Cocaine dependence, uncomplicated
§	Z20.828	Contact with and (suspected) exposure to other viral communicable diseases
§	R05.9	Cough, unspecified

Symbol	Diagnosis	
	Code	Sign or Symptom
§	E53.8	Deficiency of other specified B group vitamins
§	F32.A	Depression, unspecified
§	R19.7	Diarrhea, unspecified
	F34.81	Disruptive mood dysregulation disorder
▼	Z71.51	Drug abuse counseling and surveillance of drug abuser
§	R30.0	Dysuria
◀§	R82.5	Elevated urine level of drugs, medicaments and biological substances
§	Z03.89	Encounter for observation for other suspected diseases and conditions ruled out
§	Z01.818	Encounter for other preprocedural examination
§	Z01.89	Encounter for other specified special examinations
§	Z01.812	Encounter for preprocedural laboratory examination
§	Z11.52	Encounter for screening for COVID-19
§	Z13.1	Encounter for screening for diabetes mellitus
§	Z13.0	Encounter for screening for diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism
§	Z11.4	Encounter for screening for human immunodeficiency virus [HIV]
§	Z11.3	Encounter for screening for infections with a predominantly sexual mode of transmission
§	Z13.220	Encounter for screening for lipid disorders
★§♀	Z12.4	Encounter for screening for malignant neoplasm of cervix
★§	Z12.11	Encounter for screening for malignant neoplasm of colon
★§♂	Z12.5	Encounter for screening for malignant neoplasm of prostate
§	Z13.228	Encounter for screening for other metabolic disorders
§	Z13.29	Encounter for screening for other suspected endocrine disorder
§	Z11.59	Encounter for screening for other viral diseases
§	Z11.1	Encounter for screening for respiratory tuberculosis
§	Z13.9	Encounter for screening, unspecified
❖❖§	Z51.81	Encounter for therapeutic drug level monitoring
◀§	I10	Essential (primary) hypertension
≈§	R35.0	Frequency of micturition
	F41.1	Generalized anxiety disorder
✦♂§	B20	Human immunodeficiency virus [HIV] disease
§	R73.9	Hyperglycemia, unspecified
§	E78.5	Hyperlipidemia, unspecified
§	E03.9	Hypothyroidism, unspecified
§	R73.01	Impaired fasting glucose
	D50.9	Iron deficiency anemia, unspecified
Δ★	Z79.01	Long term (current) use of anticoagulants

Symbol	Diagnosis	
	Code	Sign or Symptom
Δ⊕§	Z79.891	Long term (current) use of opiate analgesic
§	M54.5Ø	Low back pain, unspecified
§	F33.2	Major depressive disorder, recurrent severe without psychotic features
§	F33.Ø	Major depressive disorder, recurrent, mild
§	F33.1	Major depressive disorder, recurrent, moderate
§	F33.3	Major depressive disorder, recurrent, severe with psychotic symptoms
§	F33.9	Major depressive disorder, recurrent, unspecified
§	F32.1	Major depressive disorder, single episode, moderate
§	F32.9	Major depressive disorder, single episode, unspecified
§	E78.2	Mixed hyperlipidemia
▲§	D7Ø.9	Neutropenia, unspecified
§	F17.2ØØ	Nicotine dependence, unspecified, uncomplicated
⊙§	E66.9	Obesity, unspecified
§	F11.11	Opioid abuse, in remission
§	F11.1Ø	Opioid abuse, uncomplicated
§	F11.24	Opioid dependence with opioid-induced mood disorder
§	F11.288	Opioid dependence with other opioid-induced disorder
§	F11.23	Opioid dependence with withdrawal
§	F11.21	Opioid dependence, in remission
§	F11.2Ø	Opioid dependence, uncomplicated
§	F11.9Ø	Opioid use, unspecified, uncomplicated
§	R73.Ø9	Other abnormal glucose
§	F31.89	Other bipolar disorder
►§	G89.29	Other chronic pain
§	R53.83	Other fatigue
	R68.89	Other general symptoms and signs
§	M51.369	Other intervertebral disc degeneration, lumbar region without mention of lumbar back pain or lower extremity pain
§	M51.26	Other intervertebral disc displacement, lumbar region
Δ⊕	Z79.899	Other long term (current) drug therapy
§	R53.81	Other malaise
▲§	D7Ø.8	Other neutropenia
§	F19.12Ø	Other psychoactive substance abuse with intoxication, uncomplicated
§	F19.1Ø	Other psychoactive substance abuse, uncomplicated
§	F19.2Ø	Other psychoactive substance dependence, uncomplicated
◀§	R79.89	Other specified abnormal findings of blood chemistry
§	F5Ø.89	Other specified eating disorder
	M47.896	Other spondylosis, lumbar region
§	F15.1Ø	Other stimulant abuse, uncomplicated
§	F15.21	Other stimulant dependence, in remission
§	F15.2Ø	Other stimulant dependence, uncomplicated
	M25.561	Pain in right knee
	M25.5Ø	Pain in unspecified joint
§	R52	Pain, unspecified
§	F41.Ø	Panic disorder [episodic paroxysmal anxiety]
◀§	F2Ø.Ø	Paranoid schizophrenia
	Z76.89	Persons encountering health services in other specified circumstances
	M96.1	Postlaminectomy syndrome, not elsewhere classified
	F43.12	Post-traumatic stress disorder, chronic
	F43.1Ø	Post-traumatic stress disorder, unspecified
§	R73.Ø3	Prediabetes
§	E78.ØØ	Pure hypercholesterolemia, unspecified
§	M54.12	Radiculopathy, cervical region

Symbol	Diagnosis	
	Code	Sign or Symptom
§	M54.16	Radiculopathy, lumbar region
§	M54.17	Radiculopathy, lumbosacral region
	M46.1	Sacroiliitis, not elsewhere classified
§	F25.Ø	Schizoaffective disorder, bipolar type
§	F25.1	Schizoaffective disorder, depressive type
§	F25.9	Schizoaffective disorder, unspecified
◀§	F2Ø.9	Schizophrenia, unspecified
§	F13.2Ø	Sedative, hypnotic or anxiolytic dependence, uncomplicated
	M48.Ø62	Spinal stenosis, lumbar region with neurogenic claudication
	M47.812	Spondylosis without myelopathy or radiculopathy, cervical region
	M47.816	Spondylosis without myelopathy or radiculopathy, lumbar region
	M47.817	Spondylosis without myelopathy or radiculopathy, lumbosacral region
□§♂	E29.1	Testicular hypofunction
◆○§	E11.22	Type 2 diabetes mellitus with diabetic chronic kidney disease
◆§	E11.65	Type 2 diabetes mellitus with hyperglycemia
◆‡§	E11.69	Type 2 diabetes mellitus with other specified complication
◆§	E11.9	Type 2 diabetes mellitus without complications
◀§	F2Ø.3	Undifferentiated schizophrenia
§	R1Ø.9	Unspecified abdominal pain
◀§	R82.9Ø	Unspecified abnormal findings in urine
	F39	Unspecified mood [affective] disorder
§	F29	Unspecified psychosis not due to a substance or known physiological condition
‡§	B19.2Ø	Unspecified viral hepatitis C without hepatic coma
▼§	N39.Ø	Urinary tract infection, site not specified
§	M54.51	Vertebrogenic low back pain
§	E55.9	Vitamin D deficiency, unspecified

Legend for Symbols

- ◀ Use additional code to identify any retained foreign body, if applicable (Z18.-)
- § Excludes1 (mutually exclusive diagnosis code) notation applies (please refer to the ICD-10-CM manual)
- ☆ Use additional code, where applicable, to identify:
 - exposure to environmental tobacco smoke (Z77.22)
 - exposure to tobacco smoke in the perinatal period (P96.81)
 - history of tobacco dependence (Z87.891)
 - occupational exposure to environmental tobacco smoke (Z57.31)
 - tobacco dependence (F17.-)
 - tobacco use (Z72.0)
- ▲ Use additional code to identify:
 - major osseous defect, if applicable (M89.7-)
 - personal history of (healed) osteoporosis fracture, if applicable (Z87.310)
- Use additional code for blood alcohol level, if applicable (Y90.-)
- Code also any associated medical condition and intellectual disabilities
- ▶ Code also related psychological factors associated with pain (F45.42)
- ✦ Use additional code to identify resistance to antimicrobial drugs (Z16.-)
- ✓ Use additional code for drug abuse or dependence (F11-F16, F18-F19)
- ★ Use additional code to identify any family history of malignant neoplasm (Z80.-)
- ♀ Female diagnosis only
- ♂ Male diagnosis only
- ❖ Code also condition requiring care
- ◇ Code also any long-term (current) drug therapy (Z79.-)
- ◀ Use additional code, if applicable, to identify:
 - other specified cognitive deficit (R41.84-)
- ≈ Code first, if applicable, any causal condition, such as:
 - enlarged prostate (N40.1)
- ◇ Code first Human immunodeficiency virus [HIV] disease complicating pregnancy, childbirth and the puerperium, if applicable (O98.7-)
Use additional code(s) to identify all manifestations of HIV infection
- △ Code also any follow-up examination (Z08-Z09)
- ⊕ Code also any therapeutic drug level monitoring (Z51.81)
- ▲ Use additional code for any associated:
 - fever (R50.81)Code also, if applicable, mucositis (J34.81, K12.3-, K92.81, N76.81)
- ⊙ Code first obesity complicating pregnancy, childbirth and the puerperium, if applicable (O99.21-)
Use additional code to identify body mass index (BMI), if known, for adults (Z68.1-Z68.45) or pediatrics (Z68.5-)
- Use additional code for adverse effect, if applicable, to identify drug (T36-T50 with fifth or sixth character 5)
- ◆ Use additional code to identify control using:
 - insulin (Z79.4)
 - injectable non-insulin antidiabetic drugs (Z79.85)
 - oral antidiabetic drugs (Z79.84)
 - oral hypoglycemic drugs (Z79.84)
- Use additional code to identify stage of chronic kidney disease (N18-1-N18.6)
- ‡ Use additional code to identify complication
- ▼ Use additional code (B95-B97) to identify infectious agent

